

Temporo-mandibular disorders are an important comorbidity of migraine and may be clinically difficult to distinguish them from tension-type headache

Desordens temporomandibulares são comorbidade importante da migrânea e podem ser clinicamente difíceis de distinguir da cefaleia tipo tensional

Ariovaldo Alberto da Silva Júnior¹, Karina Viana Brandão², Bruno Engler Faleiros³, Rafael Mattos Tavares³, Rodrigo Pinto Lara³, Eduardo Januzzi⁴, Anísio Bueno de Carvalho⁴, Eliane Maria Duarte de Carvalho⁴, João Bosco Lima Gomes⁵, Frederico Mota Gonçalves Leite⁵, Betania Mara Franco Alves⁶, Rodrigo Santiago Gómez⁷, Antônio Lúcio Teixeira⁸

ABSTRACT

Clinical differentiation between the primary headaches and temporomandibular disorders (TMD) can be challenging. **Objectives:** To investigate the relationship between TMD and primary headaches by conducting face to face assessments in patients from an orofacial pain clinic and a headache tertiary center. **Method:** Sample consists of 289 individuals consecutively identified at a headache center and 78 individuals seen in an orofacial pain clinic because of symptoms suggestive of TMD. **Results:** Migraine was diagnosed in 79.8% of headache sufferers, in headache tertiary center, and 25.6% of those in orofacial pain clinic ($p < 0.001$). Tension-type headache was present in 20.4% and 46.1%, while the TMD painful occurred in 48.1% and 70.5% respectively ($p < 0.001$). **Conclusion:** TMD is an important comorbidity of migraine and difficult to distinguish clinically from tension-type headache, and this headache was more frequent in the dental center than at the medical center.

Keywords: primary headache, temporomandibular disorders, migraine.

RESUMO

A diferenciação clínica entre as cefaleias primárias e as disfunções temporomandibulares (DTM) pode ser desafiadora. **Objetivos:** Investigar a relação entre DTM e cefaleias primárias conduzindo uma avaliação face a face entre pacientes de um centro de dor orofacial e de um centro terciário de cefaleia. **Método:** A amostra consistiu de 289 indivíduos avaliados consecutivamente em um centro terciário de cefaleia e 78 indivíduos de uma clínica orofacial. **Resultados:** A migrânea foi diagnosticada em 79,8% dos pacientes do centro de cefaleia e 25,6% dos pacientes do centro de dor orofacial. A cefaleia do tipo tensional esteve presente em 20,4% e 46,1%, enquanto as DTM dolorosas ocorreram em 48,1% e 70,5% respectivamente ($p < 0,001$). **Conclusão:** DTM é uma comorbidade importante da migrânea e difícil de distinguir clinicamente da cefaleia do tipo tensional, tanto que esta cefaleia foi mais frequente no centro odontológico do que no centro médico.

Palavras-chave: cefaleia primaria, desordens temporomandibular, migrânea.

Headache is a frequent cause of medical consultation at all levels of care. In the general population, headaches affect nearly 50% of the subjects¹. In Brazil, migraine affects around 10% of the adults, while tension-type headache (TTH)

Serviço de Neurologia, Hospital das Clínicas, Universidade Federal de Minas Gerais, Belo Horizonte MG, Brazil.
Orofacial Pain Clinic- Unifenas-BH, Belo Horizonte MG, Brazil

¹Attending Neurologist, Serviço de Neurologia, Hospital das Clínicas, Universidade Federal de Minas Gerais, and Professor of Neurology, Faculdade de Medicina, Unifenas-BH, Belo Horizonte MG, Brazil;

²Medical student, Faculdade de Medicina, Unifenas-BH, Belo Horizonte MG, Brazil;

³Medical student, Faculdade de Medicina, Universidade Federal de Minas Gerais, Belo Horizonte MG, Brazil;

⁴Attending Dentist, Orofacial Pain Clinic-Unifenas-BH, Belo Horizonte MG, Brazil;

⁵Attending Dentist, Serviço de Neurologia, Hospital das Clínicas, Universidade Federal de Minas Gerais, Belo Horizonte MG, Brazil;

⁶Attending Physiotherapist, Serviço de Neurologia, Hospital das Clínicas, Universidade Federal de Minas Gerais, Belo Horizonte MG, Brazil;

⁷Attending Neurologist, Departamento de Clínica Médica, Faculdade de Medicina, Universidade Federal de Minas Gerais, Belo Horizonte MG, Brazil;

⁸Professor of Neurology, Departamento de Clínica Médica, Faculdade de Medicina Universidade Federal de Minas Gerais, Belo Horizonte MG, Brazil.

Correspondence: Ariovaldo Alberto da Silva Jr. Núcleo de Apoio à Pesquisa, Unifenas-BH, Rua Libano, 66 / Bairro Itapoã; 31710-030 Belo Horizonte MG - Brasil.

Support: Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG).

Conflict of interest: There is no conflict of interest to declare.

Received 09 February 2013; Received in final form 26 September 2013; Accepted 03 October 2013.

happens in other 38%^{1,2}. In a face-to-face study in which the entire population of a city in Brazil was investigated, one-year prevalence of headache was 65.4%³, and distribution of headache types were in alignment with other studies from Brazil^{4,5}. Headaches are a frequent cause of consultation in the specialty care^{6,2}.

The temporomandibular disorders (TMD) is a collective term that includes many clinic problems that involve the masticatory muscles, the temporomandibular joint (TMJ) and associated structures or both. The most common symptom is pain, and it can be aggravated by chewing or other jaw function, and can also restrict the function related to the associated structures, and articular sounds are frequent. The prevalence of these symptoms can affect until 75% of the adult population and it is estimated that 40% to 60% of general population present any signs or symptoms of TMD⁷. Although the prevalence of TMD varies widely, in the Brazilian general population over 40% of subjects have at least one TMD symptom, and 10% have three or more⁸. TMJ pain affects from 5-16% of the general population and, as for migraine headaches, women are disproportionately affected⁸⁻¹⁰.

The myofascial pain (MP) is one of the most common TMD and it consists in a painful condition characterized by pain in local areas in the jaw, temples, face, pre auricular area, or inside the ear with the presence of trigger-points (TP). These (TP) can produce a characteristic pattern of irradiated pain or autonomic symptoms when stimulated⁷.

There are many diagnostic systems for TMD of which MP is a part. The Research Diagnostic Criteria for Temporomandibular Disorders (RDC/TMD) was created to provide classification criteria for the condition that are universally accepted and validated. A diagnosis of MP by the RDC/TMD requires pain to be reported by the subject in response to palpation of three or more sites of the masticatory muscles¹¹.

Specific primary headache syndromes, in special migraine, and TMD are comorbid, as demonstrated by clinic-based and population studies. Additionally, TMD is a risk factor for increased headache frequency^{10,12-20}. Nonetheless, as with many of the identified risk factors for increased migraine frequency, most of the studies diagnosed both migraine and TMD based on self-reported symptoms or on questionnaires. Accordingly, the aim of this study was to investigate the relationship between TMD and headache syndromes by conducting rigorous face to face assessments in patients from an orofacial pain clinic and a headache tertiary center.

METHOD

Participants of this study consist of 289 individuals consecutively identified at the Headache Specialty Center at *Universidade Federal de Minas Gerais* (Belo Horizonte,

Brazil), as well as 78 individuals who sought care with complaints suggestive of TMD at the Odontologic Training and Specializing Center (Belo Horizonte, Brazil). All patients were independently assessed by neurologists with experience in headache medicine. Headaches were diagnosed according to the Second Edition of the International Classification of Headache Disorders (ICHD-2)²¹ after a structured interview and clinical examination. Headaches of interest included migraine, TTH, chronic daily headache (CDH), and medication overuse headache, which was diagnosed only after detoxification, as per the ICHD-2.

Oral and muscular assessments were respectively conducted by dentists and physical therapists. Trigger points, threshold for pain and the temporomandibular joint were rigorously assessed. TMD was assessed using the RDC/TMD²² through an interview and, subsequently, a physical examination. The RDC/TMD consists of a dual-axis approach (Axis I and II), composed by a questionnaire and a physical examination made by dentists and physiotherapists previously calibrated by a gold standard examiner, who provided an overview of all procedures and helped them to practice on each other and other subjects recording and comparing the results from the same subjects evaluated according to the International RDC/TMD Consortium specifications. Additionally, the head and neck muscles were examined and pain sensitivity to 2.0 kg straight pressure was assessed using a Likert pain scale. Examiners were blinded to each other diagnoses.

Data were summarized using summary tables and descriptive statistics. Non-parametric data between groups were compared using the Chi-Squared test or the Fischer test (when anticipated values were small). For continuous variables, the Mann-Whitney test was used. Gaussian variables were contrasted using the T-test.

Protocol and all forms were reviewed and approved by the Investigation Review Board of the Federal University of Minas Gerais, and by the Committee (0500.203.000-10) Research Ethics Unifenas, Belo Horizonte, Brazil (217/2010).

RESULTS

Among individuals with headache (n=289), mean age was 42.7 years, while in individuals with TMD symptoms, it was 37.1 years. Women represented 86.9% of headache sufferers and 75.6% in the TMDs group (p=0.015). Most participants had less than 8 years of formal education. Demographic data are presented in Table 1.

Migraine was diagnosed in 79.8% of headache sufferers and 25.6% of those with TMDs (p<0.001). TTH happened in 20.4% and 46.1% respectively (p<0.001). CDH with medication overuse was the diagnosed in 16.6% of headache sufferers and 3.8% of TMD sufferers (p=0.004) (Table 2).

Table 1. Demographic data at the two specialty centers

	Headache center (n=289)	Oral clinic (n=78)	p-value
Sex			
Women	86.9% (251)	75.6% (59)	0.015**
Men	13.1% (38)	24.4% (19)	
Years of school			
< 8 years	52.3% (136)	23.2% (16)	<0.001*
8-15 years	35.0% (91)	59.4% (41)	
> 15 years	12.7% (33)	17.4% (12)	

(*) Pearson's Chi-square Test; (**) Mann-Whitney Test.

TMD happened in 67.1% of those with headache and 97.4% of those with TMDs. Among the TMD sufferers, painful muscular dysfunction happened in 49.5% of those with headache vs. 23.1% in those from the TMDs group ($p<0.001$). Articular TMD was rarely observed among those from the headache group (1.4%) but happened in 15.4% in those from the TMDs group ($p<0.001$). Coexisting muscular and articular TMD happened in 59.0% of the TMDs patients (Table 3). TMD with myofascial pain happened in 48.1% of headache sufferers and 70.5% of those with TMDs ($p<0.001$). Local

muscle soreness was more common in the headache sufferers than in those with TMDs ($p=0.047$) (Table 4).

DISCUSSION

Headache and TMD are two painful disorders that are comorbid and may sometimes be disabled, requiring specialized treatment. Better characterization of the comorbidity is necessary as a prelude to customized treatment. Our study adds to the field by formally assessing patients from two specialty clinics using gold-standard diagnostic criteria.

As reported by others, women were more commonly affected by both disorders, specially at young adulthood and middle-age^{1,3-6,8-10,12-14,22}, and it seems that the female predominance is amplified at the specialty care, relative to community studies^{3,6,20,23}. Since migraine and TMD have different physiopathology, we did not expect to find identical gender ratio in both groups. Nonetheless, part of this difference may be justified by the fact that when TMD and migraine co-occur, migraine tends to be more severe than when they do not, which would channel women with more severe TMD to the

Table 2. Headache types and frequency as a function of place of enrollment.

	Headache center (n=289)		Oral clinic (n=78)		p-value
	%	95%CI	%	95%CI	
Migraine	79.89	[74.16; 83.63]	25.64	[15.73; 35.55]	<0.001*
Tension-type headache	20.42	[15.74; 25.09]	46.15	[34.84; 57.47]	<0.001*
Chronic daily headache	16.61	[12.29; 20.93]	3.85	[0.52; 8.21]	0.004*

(*) Pearson's Chi-Square Test.

Table 3. Temporomandibular diagnosis as a function of place of enrollment.

	Headache center (n=289)	Oral clinic (n=78)	p-value
No TDM	32.9% (95)	2.6% (2)	<0.001*
Muscular	49.5% (143)	23.1% (18)	
Articular	1.4% (4)	15.4% (12)	
Muscular and articular	16.3% (47)	59.0% (46)	

TDM: temporomandibular disorders; (*) Pearson's Chi-Square test;

Table 4. Temporomandibular disorder types as function of place of enrollment.

	Headache center		Oral clinic		p-value
	Prevalence	95%CI	Prevalence	95%CI	
Disc dislocation with reduction	3.46%	[1.34; 5.58]	28.21%	[17.99; 38.42]	<0.001*
Disc dislocation without reduction	0.35%	[0.34; 1.03]	10.26%	[3.37; 17.14]	<0.001*
Spontaneous dislocation	0.69%	[0.27; 1.65]	7.69%	[1.65; 13.74]	0.002**
Synovitis/ capsulitis	17.65%	[9.24; 26.06]	42.31%	[31.10; 53.52]	<0.001*
Myofascial pain	48.10%	[42.30; 53.89]	70.51%	[60.17; 80.86]	<0.001*
Local muscle soreness	18.34%	[13.85; 22.83]	8.97%	[2.49; 15.46]	0.047*
Myospasm	0%	-	8.97%	[2.49; 15.46]	<0.001**

(*) Pearson's Chi-Square test; (**) Fisher's exact test.

headache center rather to the TMD center¹⁰. Furthermore, it is well established that women are more likely than men to suffer pain episodes as a function of neuroendocrine events and of the reproductive stage^{10,24-26}. Menstrual migraine is well known to be more severe and refractory to treatment than non-menstrual migraine^{10,24}, and a limitation of our study was not to characterize this migraine subgroup, since this would require longitudinal follow-up and use of dairies.

We found strong association between migraine and TMD pain, as previously described^{10,12-20}. Gonçalves et al., in a population-based study, found that TMD symptoms are more likely to occur among those who present any headache type. When three or more TMD symptoms were present, headache occurred in 72.8% versus 37.9% of those who did not present any TMD symptoms¹³, as confirmed by others¹⁵. Furthermore, when assessing a clinic-based sample, they found that the comorbidity of TMD with migraine, CDH, and TTH, happened only for the muscular form, and not for the articular type¹⁸. They also found that severity of TMD correlated with increased headache frequency^{15,18}. Our data strongly support their findings and this is of importance, since Bevilacqua-Grossi et al. reported that TMD with myofascial pain is strongly associated to increased headache frequency²⁷.

Of interest is the fact that we found more TTH sufferers in the TMD clinic than in the headache center. TTH is

the most prevalent of the primary headaches in the general population, with prevalences ranging from 38-78%^{1,28}. In a telephone-based interview conducted in Brazil, prevalence was 13%, while probable TTH happened in other 22.6%. Thus, further studies should focus on the relationship between TMD and TTH. In another clinic-based study, findings were similar to ours²⁹.

Finally, we also found evidence that CDH and TMD are comorbid, as suggested by others, where painful TMD was strongly associated with specific CDH types^{13,14,18,20,27,30}.

This study has several limitations, in addition to the lack of controlling for menstrual cycle. First, both groups come from specialty care, without a contemporary control from the population. Second, psychiatric comorbidities were not assessed and this is of importance, since they may be related to both headaches and TMD. Finally, patients were not matched by age and gender (since they were consecutively enrolled).

We conducted a study independently identifying patients from two specialty centers, one in headache and one in TMD. We found strong evidence of comorbidity between both disorders and we confirmed previous findings about the specificity of TMD pain and of TMD with muscular symptoms in the comorbidity. We also found evidence of an increased cluster of TTH sufferers in the TMD clinic, and this finding must be further elucidated.

References

1. Stovner LJ, Hagen K, Jensen R, et al. The global burden of headache: a documentation of headache prevalence and disability worldwide. *Cephalalgia* 2007;27:193-210.
2. Crystal SC, Robbins MS. Epidemiology of tension-type headache. *Curr Pain Headache Rep* 2010;14:449-454.
3. Silva-Júnior AA, Krymchantowski A, Moreira P, Vasconcelos L, Gomez RS, Teixeira AL. Prevalence of headache in the entire population of a small city in Brazil. *Headache* 2009;49:895-899.
4. Queiroz LP, Barea LM, Blank N. An epidemiological study of headache in Florianópolis, Brazil. *Cephalalgia* 2005;26:122-127.
5. Pahim LS, Menezes AMB, Lima R. Prevalence and factors associated to migraine in adult population, Southern Brazil. *Rev Saúde Pública* 2006;40:692-698.
6. Silva-Júnior AA, Faleiros BE, Santos TM, Gómez RS, Teixeira AL. Relative frequency of headache types: a longitudinal study in the tertiary care. *Arq Neuropsiquiatr* 2010;68:878-881.
7. Leite FMG, Atallah AN, El Dib RP, et al. Cyclobenzaprine for the treatment of myofascial pain in adults. *Cochrane Database of Systematic Reviews* 2009, Issue 3. Art. No.: CD006830. DOI: 10.1002/14651858.CD006830.pub3.
8. Gonçalves AGG, Fabbro ALD, Campos JADB, Bigal ME, Speciali JG. Symptoms of temporomandibular disorders in the population: an epidemiological study. *J Orofac Pain* 2010;24:270-278.
9. Mobilio N, Casetta I, Cesnik E, Catapano S. Prevalence of self-reported symptoms related to temporomandibular disorders in an Italian population. *J Oral Rehab* 2010;38:884-890.
10. Plesh O, Adams SH, Gansky SA. Temporomandibular Joint and muscle disorder-type pain and comorbid pains in a national US sample. *J Orofac Pain* 2011;25:190-198.
11. Manfredini D, Chiappe G, Bosco M. Research diagnostic criteria for temporomandibular disorders (RDC/TMD) axis I diagnoses in an Italian patient population. *J Oral Rehab* 2006;33:551-558.
12. Silva-Junior AA, Costa EC, Gomes JB, et al. Chronic headache and comorbidities: a two-phase, population-based, cross-sectional study. *Headache* 2010;50:1306-1312.
13. Gonçalves DAG, Bigal ME, Jales LCF, Camparis CM, Speciali JG. Headache and symptoms of temporomandibular disorder: an epidemiological study. *Headache* 2010;50:231-241.
14. Gonçalves DAG, Speciali JG, Jales LCF, Camparis CM, Bigal ME. Temporomandibular symptoms, migraine and chronic daily headaches in the population. *Neurology* 2009;73:645-646.
15. Ballegaard V, Thede-Schmidt-Hansen P, Svensson P, Jensen R. Are headache and temporomandibular disorders related? A blinded study. *Cephalalgia* 2008;28:832-841.
16. Fragoso YD, Alves HHC, Garcia SO, Finkelsztein A. Prevalence of parafunctional habits and temporomandibular dysfunction symptoms in patients attending a tertiary headache clinic. *Arq Neuropsiquiatr* 2010;68:377-380.
17. Kang JK, Ryu JW, Choi JH, Merrill RL, Kim ST. Application of ICDH-II criteria for headaches in a TMJ and orofacial pain clinic. *Cephalalgia* 2009;30:37-41.
18. Gonçalves DAG, Camparis CM, Speciali JG, Franco AL, Castanharo SM, Bigal ME. Temporomandibular disorders are differentially associated with headache diagnoses: a controlled study. *Clin J Pain* 2011;27:611-615.
19. Özkan NÇ, Özkan F. The relationship of temporomandibular disorders with headaches: a retrospective analysis. *Agri* 2011;23:13-17.
20. Franco AL, Gonçalves DAG, Castanharo SM, Speciali JG, Bigal ME, Camparis CM. Migraine is the most prevalent primary headache

- in individuals with temporomandibular disorders. *J Orofac Pain* 2010;24:287-292.
21. Headache Classification Committee. The international classification of headache disorders, cranial neuralgia and facial pain. *Cephalalgia* 2004;2(Suppl 1):S1-S160.
 22. Dworkin SF, Sherman J, Mancl L, Ohrbach R, LeResche L, Truelove E. Reliability, validity, and clinical utility of the research diagnostic criteria for temporomandibular disorders axis II scales: depression, non-specific physical symptoms, and graded chronic pain. *J Orofac Pain* 2002;16:207-220.
 23. Felício AC, Bichuetti DB, Santos WA, Godeiro Junior CO, Marin LF, Carvalho DS. Epidemiology of primary and secondary headaches in a Brazilian tertiary-care center. *Arq Neuropsiquiatr* 2006;64:41-46.
 24. Nappi RE, Berga SL. Migraine and reproductive life. *Handb Clin Neurol* 2010;97:303-322.
 25. Liverman CS, Brown JW, Sandhir R, McCarson KE, Berman NE. Role of the oestrogen receptors GPR30 and ERalpha in peripheral sensitization: relevance to trigeminal pain disorders in women. *Cephalalgia* 2009;29:729-741.
 26. Gupta S, McCarson KE, Welch KM, Berman NE. Mechanisms of pain modulation by sex hormones in migraine. *Headache* 2011;51:905-922.
 27. Bevilacqua-Grossi D, Lipton RB, Napchan U, Grosberg B, Ashina S, Bigal ME. Temporomandibular disorders and cutaneous allodynia are associated in individuals with migraine. *Cephalalgia* 2010;30:425-432.
 28. Queiroz LP, Peres MF, Piovesan EJ, et al. A nationwide population-based study of tension-type headache in Brazil. *Headache* 2009;49:71-78.
 29. Koenig MA, Gladstein J, McCarter RJ, Hershey AD, Wasiewski W, Pediatric Committee of the American Headache Society. Chronic daily headache in children and adolescents presenting to tertiary headache clinics. *Headache* 2002;42:491-500.
 30. Stuginski-Barbosa J, Macedo HR, Bigal ME, Speciali JG. Signs of temporomandibular disorders in migraine patients: a prospective, controlled study. *Clin J Pain* 2010;26:418-421.